

Bayer Healthcare LLC.

- PE Product Labels –



FINISH AREA



Alka-Seltzer PLUS

FREE VARN 841955881



84195588

Do not use if pouch is torn or open

36 EFFERVESCENT TABLETS

ORIGINAL

Alka-Seltzer PLUS

COLD FORMULA



Drug Facts

Active ingredients (in each tablet)

Aspirin 325 mg (NSAID)*.....Pain reliever/fever reducer

Chlorpheniramine maleate 2 mg.....Antihistamine

Phenylephrine bitartrate 7.8 mg.....Nasal decongestant

*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - headache
 - runny nose
 - nasal and sinus congestion
 - sneezing
 - sore throat
 - temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have
 - asthma
 - diabetes
 - thyroid disease
 - glaucoma
 - difficulty in urination due to enlargement of the prostate gland
 - a breathing problem such as emphysema or chronic bronchitis
 - a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for
 - gout
 - diabetes
 - arthritis
- taking sedatives or tranquilizers

Drug Facts (continued)

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children
- you may get drowsy
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 4 hours. Do not exceed 8 tablets in 24 hours or as directed by a doctor.
- children under 12 years: do not use

Alka-Seltzer PLUS

Aspirin (NSAID) / Pain reliever- fever reducer • Chlorpheniramine maleate / Antihistamine • Phenylephrine bitartrate / Nasal decongestant

COLD FORMULA

SPARKLING ORIGINAL

Nasal Congestion

Runny Nose

Headache & Body Ache

Sore Throat

Sinus Pressure

Dist. by: Bayer HealthCare LLC
Whippany, NJ 07981

Bayer, the Bayer Cross and Alka-Seltzer Plus are registered trademarks of Bayer.

Expiry Date:

Lot:

Drug Facts (continued)

Other information

- each tablet contains: sodium 474 mg
- Phenylephrine: Contains Phenylephrine 8.4 mg Per Tablet
- store at room temperature. Avoid excessive heat.

Inactive ingredients acesulfame potassium, anhydrous citric acid, aspartame, calcium silicate, dimethylpolysiloxane, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

Questions or comments? 1-800-886-0369 (Mon-Fri 9 AM - 5 PM EST)

MADE IN MEXICO

FPO 100% UPC

16500 53765 6

Bayer









C3100

[illegible]

[illegible]

RESTRICTED







Do not take these products at the same time.

Alka-Seltzer Plus® Maximum Strength Severe Sinus Congestion & Cough Day Liquid Gels

Drug Facts

Active ingredients (in each capsule)

Acetaminophen 325 mg..... Pain reliever/fever reducer
 Dextromethorphan hydrobromide 10 mg..... Cough suppressant
 Phenylephrine hydrochloride 5 mg..... Nasal decongestant

Purposes

Alka-Seltzer Plus® Maximum Strength Severe Sinus Congestion & Cough Night Liquid Gels

Drug Facts

Active ingredients (in each capsule)

Acetaminophen 325 mg..... Pain reliever/fever reducer
 Dextromethorphan hydrobromide 10 mg..... Cough suppressant
 Doxylamine succinate 6.25 mg..... Antihistamine
 Phenylephrine hydrochloride 5 mg..... Nasal decongestant

Purposes

Alka-Seltzer Plus® Maximum Strength Severe Sinus Congestion & Cough Day Liquid Gels

Drug Facts (continued)

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - nasal congestion
 - sinus congestion and pressure
 - headache
 - minor aches and pains
 - cough
 - sore throat
- helps clear nasal passages and shrinks swollen membranes
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

LIFT HERE

Alka-Seltzer Plus® Maximum Strength Severe Sinus Congestion & Cough Night Liquid Gels

Drug Facts (continued)

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - nasal congestion
 - sinus congestion and pressure
 - headache
 - minor aches and pains
 - cough
 - sore throat
 - runny nose
 - sneezing
- helps clear nasal passages and shrinks swollen membranes
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:


- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

O



Cut Edge

Alka-Seltzer Plus® Maximum Strength Severe Sinus Congestion & Cough Day Liquid Gels
Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough with excessive phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Alka-Seltzer Plus® Maximum Strength Severe Sinus Congestion & Cough Night Liquid Gels
Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have


- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

INSIDE of EC Flap

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org



Alka-Seltzer Plus® Maximum Strength Severe Sinus Congestion & Cough Day Liquid Gels

Drug Facts (continued)

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use

Other information

- store at room temperature. Avoid excessive heat above 40°C (104°F).

Alka-Seltzer Plus® Maximum Strength Severe Sinus Congestion & Cough Night Liquid Gels

Drug Facts (continued)

When using this product

- do not exceed recommended dosage
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use

Alka-Seltzer Plus® Maximum Strength Severe Sinus Congestion & Cough Day Liquid Gels

Drug Facts (continued)

Inactive ingredients FD&C yellow #6, FD&C red #40, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)

Alka-Seltzer Plus® Maximum Strength Severe Sinus Congestion & Cough Night Liquid Gels

Drug Facts (continued)

Other information

- store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients FD&C blue #1, D&C yellow #10, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

Questions? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)

Made in Spain
© 2017 Bayer.
Dist. by:
Bayer HealthCare LLC
Whippany, NJ 07981
Bayer, the Bayer Cross and Alka-Seltzer Plus are registered trademarks of Bayer.

Maximum Strength Per 4 Hour Dose
Do not use if blister is cut or broken.
DOES NOT CONTAIN ASPIRIN

Bayer

LOT/EXP

LOT & EXP CODE AREA

85740969

8 1/2

3/4

1/2

8 1/2





7312

Drug Facts

Active ingredients (in each capsule)

Acetaminophen 325mg Pain reliever/fever reducer

Dextromethorphan hydrobromide 10 mg Cough suppressant

Phenylephrine hydrochloride 5 mg Nasal decongestant

Purposes

- temporarily relieves these symptoms due to a cold or flu:
- nasal congestion
- sinus congestion and pressure
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Drug Facts (continued)

Ask a doctor before use if you have:

- low blood pressure
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with excessive phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- prolapsed or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the listed Warning Drug Interactions.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if:

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nausea, vomiting, or dizziness occurs

Directions

- do not take more than the recommended dose
- adults and children 12 years and over take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use

Other information

- store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients: FD&C yellow No. 6, ferric oxide, gelatin, glycerin, polyethylene glycol, potassium aluminum silicate, povidone, propylene glycol, purified water, xanthan, sodium hydroxide, sorbitol sorbitate solution, titanium dioxide

Drug Facts (continued)

Questions or comments?

1-800-486-4368 (Only for 9AM - 5PM EST)

MAXIMUM STRENGTH

Sinus Congestion & Pain

Maximum Strength Per 4 Hour Dose

POWERMAX GELS

25% SMALLER LIQUID GELS

Alka-Seltzer PLUS

MAXIMUM STRENGTH

Sinus Congestion & Pain

POWERMAX GELS

CONCENTRATED FORMULA

MAXIMUM STRENGTH

Sinus Congestion & Pain

Do not use if blister is cut or broken. DOES NOT CONTAIN ASPIRIN

POWERMAX GELS

Acetaminophen / Pain Reliever-Fever Reducer

Dextromethorphan HBr / Cough Suppressant

Phenylephrine HCl / Nasal Decongestant

PARENTS:

Learn about these medical claims

www.StupidSchools.org

LOT/EXP:

For FAST Relief Of:

- Nasal/Sinus Congestion
- Headache, Body Ache, Sore Throat
- Sinus Pressure
- Cough

24 LIQUID GELS (Liquid Filled Capsules)

***Compared to original Alka-Seltzer Plus Liquid Gels**

ACTUAL SIZE

Bayer the Bayer Cross, Alka-Seltzer Plus, and POWERMAX are registered trademarks of Bayer.

Product of Spain




Dist. by: Bayer HealthCare LLC, Whispering, NJ 07081

879-17312 © Bayer 2021 Pat. patents.bayer.com

Bayer

0 16500 58349 3





INSIDE of EC Flap

PARTIAL GLUE LIST

6264

1/2

3/4

<div style="background-color: #f96; padding: 5px; text-align: center;"> DAY NON-DROWSY </div> <p>ACETAMINOPHEN / Pain Reliever-Fever Reducer Dextromethorphan HBr / Cough Suppressant Phenylephrine HCl / Nasal Decongestant</p> <div style="display: flex; justify-content: space-between;"> <ul style="list-style-type: none"> ➤ Nasal Congestion ➤ Headache & Body Ache ➤ Cough ➤ Sore Throat ➤ Sinus Pressure <div style="text-align: center;"> <p>12</p> <p>LIQUID GELS (liquid filled capsules)</p> </div> </div>	<div style="background-color: #008000; color: white; padding: 5px; text-align: center;"> NIGHT </div> <p>ACETAMINOPHEN / Pain Reliever-Fever Reducer Dextromethorphan HBr / Cough Suppressant Doxylamine Succinate / Antihistamine Phenylephrine HCl / Nasal Decongestant</p> <div style="display: flex; justify-content: space-between;"> <ul style="list-style-type: none"> ➤ Nasal Congestion ➤ Headache & Body Ache ➤ Cough ➤ Runny Nose ➤ Sore Throat <div style="text-align: center;"> <p>8</p> <p>LIQUID GELS (liquid filled capsules)</p> </div> </div>
--	--

<p style="text-align: center;">Day Cold & Flu Liquid Gels</p> <p>Alka-Seltzer Plus® Maximum Strength</p> <p>Drug Facts (continued)</p> <p>Inactive Ingredients: FD&C blue #1, FD&C yellow #10, gelatin, glycerin, polyethylene glycol, polydioxane, propylene glycol, purified water, silica, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide</p> <p>Questions or comments? 1-800-865-0366 (Mon-Fri 9AM - 5PM EST)</p>	<p style="text-align: center;">Night Cold & Flu Liquid Gels</p> <p>Alka-Seltzer Plus® Maximum Strength</p> <p>Drug Facts (continued)</p> <p>Inactive Ingredients: FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, polydioxane, propylene glycol, purified water, silica, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide</p> <p>Questions or comments? 1-800-865-0366 (Mon-Fri 9AM - 5PM EST)</p>
--	---

<p style="text-align: center;">Day Cold & Flu Liquid Gels</p> <p>Alka-Seltzer Plus® Maximum Strength</p> <p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ● pain, cough, or nasal congestion gets worse or lasts more than 7 days ● fever gets worse or lasts more than 3 days ● new symptoms occur ● cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. ● nervousness, dizziness, or sleeplessness occurs <p>Before use:</p> <ul style="list-style-type: none"> ● If pregnant or breast-feeding, ask a health professional <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions:</p> <ul style="list-style-type: none"> ● adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours ● or as directed by a doctor ● children under 12 years: do not use <p>Other Information:</p> <ul style="list-style-type: none"> ● store at room temperature. Avoid excessive heat above 40°C (104°F). 	<p style="text-align: center;">Night Cold & Flu Liquid Gels</p> <p>Alka-Seltzer Plus® Maximum Strength</p> <p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ● pain, cough, or nasal congestion gets worse or lasts more than 7 days ● fever gets worse or lasts more than 3 days ● new symptoms occur ● cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. ● nervousness, dizziness, or sleeplessness occurs <p>Before use:</p> <ul style="list-style-type: none"> ● If pregnant or breast-feeding, ask a health professional <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions:</p> <ul style="list-style-type: none"> ● adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours ● or as directed by a doctor ● children under 12 years: do not use <p>Other Information:</p> <ul style="list-style-type: none"> ● store at room temperature. Avoid excessive heat above 40°C (104°F).
---	---

6264

3/4

LOT EXP

Bayer

Maximum Strength Per 4 Hour Dose
Do not use if blister is cut or broken.
DOES NOT CONTAIN ASPIRIN

Made in Spain
© 2017 Bayer.
Dist. by:
Bayer Healthcare LLC
Whispery, NJ 07881
Bayer, the Bayer Cross, and
Alka-Seltzer Plus are
registered trademarks
of Bayer.

86329264

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

0 16500 55531 5




OPENING INSTRUCTIONS:
"LIFT HERE"

Cut Edge

Alka-Seltzer Plus® Maximum Strength Day Cold & Flu Liquid Gels	Alka-Seltzer Plus® Maximum Strength Night Cold & Flu Liquid Gels
<p>Drug Facts (continued)</p> <p>Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:</p> <ul style="list-style-type: none"> skin redness hives facial swelling asthma (wheezing) shock <p>If a skin or general allergic reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use:</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. if you have ever had an allergic reaction to this product or any of its ingredients in children under 12 years of age <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> liver disease heart disease high blood pressure thyroid disease diabetes cough with excessive phlegm (mucus) difficulty in urination due to enlargement of the prostate gland persistent or chronic cough such as occurs with smoking 	<p>Drug Facts (continued)</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use to sedate children.</p> <p>Do not use:</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. if you have ever had an allergic reaction to this product or any of its ingredients in children under 12 years of age <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> liver disease heart disease high blood pressure thyroid disease diabetes glaucoma cough with excessive phlegm (mucus) a breathing problem such as emphysema or chronic bronchitis difficulty in urination due to enlargement of the prostate gland persistent or chronic cough such as occurs with smoking, asthma, or emphysema <p>Ask a doctor or pharmacist before use if you are:</p> <ul style="list-style-type: none"> taking the blood thinning drug warfarin

INSIDE of EC Flap







Drug Facts

Active ingredients (in each capsule)

Acetaminophen 325 mg.....	Pain reliever/ fever reducer
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine
Phenylephrine hydrochloride 5 mg.....	Nasal decongestant

Purposes

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes
- temporarily relieves these symptoms due to a cold:
 - nasal congestion
 - sinus congestion and pressure
 - headache
 - minor aches and pains
 - cough
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Drug Facts (continued)

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Do not use to sedate children.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

LIFT HERE